Web figure 4: Unified Parkinson's disease rating scale – activities of daily living

	Intervention			No Intervention				Mean Difference	Mean Difference
Study or Subgroup	Mean	SD	Total	Mean	SD	Total	Weight	IV, Fixed, 95% CI	IV, Fixed, 95% CI
4.3.1 General Physic	otherapy	v Conf	trol						
Ellis 2005	-2.1	2.8	32	-0.3	2.3	33	71.4%	-1.80 [-3.05, -0.55]	
Fisher 2008	-1.5	2.81	10	-0.9	3.9	10	12.5%	-0.60 [-3.58, 2.38]	•
Subtotal (95% CI)			42			43	83.9%	-1.62 [-2.77, -0.47]	•
Heterogeneity: Chi²=	: 0.53, df	= 1 (P	= 0.47); I ^z = 0%					
Test for overall effect	: Z= 2.76	(P = 0	0.006)						
4.3.2 Treadmill v Cor	ntrol								
Fisher 2008	0.6	3.64	10	-0.9	3.9	10	10.2%	1.50 [-1.81, 4.81]	
Subtotal (95% CI)			10			10	10.2%	1.50 [-1.81, 4.81]	
Heterogeneity: Not ap	pplicable								
Test for overall effect	:: Z = 0.89	(P = 0	0.37)						
4.3.3 Dance v Contro	ol								
Duncan 2012	-0.6	7.5	26	1.9	8.4	26	5.9%	-2.50 [-6.83, 1.83]	
Subtotal (95% CI)			26			26	5.9%	-2.50 [-6.83, 1.83]	
Heterogeneity: Not ap	pplicable								
Test for overall effect	: Z=1.13	(P = 0	0.26)						
Total (95% CI)			78			79	100.0%	-1.36 [-2.41, -0.30]	•
Heterogeneity: Chi ² =	3.87, df	= 3 (P	= 0.28); l = 229	%				140 1
Toot for everall offeet: 7 = 2.52 /B = 0.043									-10 -5 0 5 10 Favours Intervention Favours No Intervention
Test for subgroup dif	fferences	: Chi²:	= 3.34.	df = 2 (P)	= 0.19	$3), \mathbf{f} = 4$	40.1%		rayours intervention Favours No Intervention